



Recomendaciones de la Academia Estadounidense de Cirujanos Ortopédicos

Antibióticos después del remplazo de articulación

El remplazo completo de una articulación infectada puede ser muy grave y el tratamiento puede ser muy costoso. Debido a esto, la Academia estadounidense de Cirujanos Ortopédicos recomienda que los clínicos consideren tratar con antibióticos a todos los pacientes que se someten a remplazos completos de articulación antes de que se sometan a cualquier procedimiento que pudiera llevar bacterias dentro de la corriente sanguínea. Esta tabla muestra sus recomendaciones de antibióticos profilácticos 2009. **Por favor, muestre esta tabla a su médico antes de someterse a cualquier procedimiento médico o dental.**

Si su cirujano, proveedor de atención primaria o dentista tienen alguna pregunta acerca de los antibióticos profilácticos, por favor pídale que llamen al Centro de Cirugía de Huesos y Articulaciones, al 206-598-4288 en el Centro Médico de la Universidad de Washington.

Procedure	Antimicrobial Agent	Dose	Timing	Duration
Dental	Cephalexin, cephradine, amoxicillin	2 gm by mouth	1 hour before procedure	Discontinued within 24 hours of the procedure. For most outpatient/ office-based procedures, a single pre-procedure dose is adequate.
Ophthalmic (eyes)	Gentamicin, tobramycin, ciprofloxacin, gatifloxacin, levofloxacin, moxifloxacin, ofloxacin, OR meomycin-gramicidin-polymyxin B cefazolin	Multiple eye drops topically over 2 to 24 hours OR 100 mg subconjunctivally	Consult with ophthalmologist or pharmacist for dosing regimen	
Orthopaedic (bones and joints)	Cefazolin, Cefuroxime, OR Vancomycin	1 to 2 g IV 1.5 g IV 1 g IV	Begin dose 1 hour before procedure	
Vascular (blood vessels)	Cefazolin OR Vancomycin	1 to 2 g IV 1 g IV	Begin dose 1 hour before procedure	
Esophageal, Gastroduodenal (food tube, stomach, 1st part of small intestine)	Cefazolin	1 to 2 g IV	Begin dose 1 hour before procedure	
Biliary tract (gallbladder, bile ducts)	Cefazolin	1 to 2 g IV	Consult with health care provider and/or pharmacist	
Colorectal (colon, rectum)	Neomycin + erythromycin base (oral)	1 g	Depends on time of procedure – consult with gastroenterologist and/or pharmacist	
	OR metronidazole (oral)	1 g		
Head and neck	Clindamycin + gentamicin OR cefazolin	600 to 900 mg IV 1.5 mg/kg IV 1 to 2 g IV	Begin dose 1 hour before procedure	
Obstetric and Gynecological (female organs)	Cefoxitin, cefazolin Ampicillin/sulbactam	1 to 2 g IV 3 g IV	Begin dose 1 hour before procedure	
Genitourinary (genital or urinary tract)	Ciprofloxacin	500 mg by mouth or 400 mg IV	Begin dose 1 hour before procedure	

American Academy of Orthopaedic Surgeons Recommendations

Antibiotics after joint replacement

An infected total joint replacement can be very serious, and treating it can be very costly. Because of this, the American Academy of Orthopaedic Surgeons recommends that clinicians consider treating all patients who have total joint replacements with antibiotics before they have any procedure that may bring bacteria into their bloodstream. This table shows their 2009 prophylactic antibiotic recommendations. **Please show this table to your doctor before you have any dental or medical procedures done.**

If your surgeon, primary care provider, or dentist has any questions about prophylactic antibiotics, please ask them to call the Bone and Joint Surgery Center at 206-598-4288 at University of Washington Medical Center.

Procedure	Antimicrobial Agent	Dose	Timing	Duration
Dental	Cephalexin, cephradine, amoxicillin	2 gm by mouth	1 hour before procedure	Discontinued within 24 hours of the procedure. For most outpatient/ office-based procedures, a single pre-procedure dose is adequate.
Ophthalmic (eyes)	Gentamicin, tobramycin, ciprofloxacin, gatifloxacin, levofloxacin, moxifloxacin, ofloxacin, OR meomycin-gramicidin-polymyxin B cefazolin	Multiple eye drops topically over 2 to 24 hours OR 100 mg subconjunctivally	Consult with ophthalmologist or pharmacist for dosing regimen	
Orthopaedic (bones and joints)	Cefazolin, Cefuroxime, OR Vancomycin	1 to 2 g IV 1.5 g IV 1 g IV	Begin dose 1 hour before procedure	
Vascular (blood vessels)	Cefazolin OR Vancomycin	1 to 2 g IV 1 g IV	Begin dose 1 hour before procedure	
Esophageal, Gastroduodenal (food tube, stomach, 1st part of small intestine)	Cefazolin	1 to 2 g IV	Begin dose 1 hour before procedure	
Biliary tract (gallbladder, bile ducts)	Cefazolin	1 to 2 g IV	Consult with health care provider and/or pharmacist	
Colorectal (colon, rectum)	Neomycin + erythromycin base (oral)	1 g	Depends on time of procedure – consult with gastroenterologist and/or pharmacist	
	OR metronidazole (oral)	1 g		
Head and neck	Clindamycin + gentamicin OR cefazolin	600 to 900 mg IV 1.5 mg/kg IV 1 to 2 g IV	Begin dose 1 hour before procedure	
Obstetric and Gynecological (female organs)	Cefoxitin, cefazolin Ampicillin/sulbactam	1 to 2 g IV 3 g IV	Begin dose 1 hour before procedure	
Genitourinary (genital or urinary tract)	Ciprofloxacin	500 mg by mouth or 400 mg IV	Begin dose 1 hour before procedure	