



UW Medicine

የአንጀት ምርመራ (ኮሎኖስኮፒ) ሙቪፕሬፕ(ሆድዕቃ ማፅጃ)ን በመጠቀም መዘጋጀት

የእርስዎ የኮሎኖስኮፒ (አንጀት ምርመራ) ለ \_\_\_\_\_, በ \_\_\_\_\_ ተቀጥረዋል።

| 7 ቀናት ቀደም ብሎ  | 5 ቀናት ቀደም ብሎ   | 1 ቀን ቀደም ብሎ   | በምርመራው ቀን /ዕለት   |
|---|--|---|--|
| <p>ቀን</p> <ul style="list-style-type: none"> <li><b>የደም ማቅጠኛ መድሃኒት</b> እንደ ኮሞዲን(ቀርፍሪን) ወይም ፕላቪክስ (ክሎፒዶግሬል) እየወሰዱ ከሆነ፡-             <ul style="list-style-type: none"> <li>- እነኝህን መድሃኒቶች ላዘዘው ዶክተር ይደውሉ</li> <li>- የአንጀት ምርመራ (ኮሎኖስኮፒ) እንዳለዎት እና ምርመራውን መቻሉን እንደሚያደርጉ ይግለፁለ(ላ)ት</li> <li>- ለደም ማቅጠኛ መድሃኒት የተለየ መመሪያ ካለ ይጠይቁት</li> </ul> </li> <li><b>ቀጠሮን መሰረዝ</b>:- የቀጠሮ ቀንዎን መቀየር አለብዎ መሰረዝ ከፊለን በ206 _____ ይደውሉ።</li> <li><b>ከሆስፒታል ስንብት እና መጓጓዣ</b> <ul style="list-style-type: none"> <li>- ወደቤትዎ ለመመለስ ዝግጁ ሲሆኑ አዋቂ የቤተሰብ አባል ወይም ጓደኛ መሆን አለበት የሚመልስዎት። ምርመራውን ካጠናቀቁ በኋላ ብቻዎትን ማሽከርከር፣ ታክስ ወይም አዎቶብስ አለብዎ የሆኑትን ወይም የአካል ጉዳተኛ መጓጓዣን መጠቀም አይችሉም።</li> <li>- ወደቤት የሚመልሱት አዋቂ የቤተሰብ አባል ወይም ጓደኛ አብሮዎት ካልመጣ የምርመራ ቀጠሮ የሚሰረዝ ይሆናል።</li> <li>- በቀሪዎቹ የቀንዎ ሰዓታትም ቤት ውስጥ አብሮዎት የሚሆን ኃላፊነት የሚሰማው አዋቂ ሰው እንዲኖር አጥብቀን እንመክራለን።</li> </ul> </li> </ul> | <p>ቀን</p> <ul style="list-style-type: none"> <li>ለውዝ ወይም ፍሬ ያላቸውን ምግቦች ከመመገብ ይቆጠቡ</li> <li>ብረት(አይረን) አለብዎ ብረትን የያዘ ብዝሃ-ቫይታሚኖችን ከመውሰድ ይቆጠቡ።</li> <li>የታዘዘውን ሙቪፕሬፕን (የሆድዕቃ) ማፅጃ ይውሰዱ።</li> <li>ይህንን መመሪያ በጥንቃቄ ያንብቡ</li> <li><b>የስኳር ህመም ካለበት</b> <ul style="list-style-type: none"> <li>- የስኳር መድሃኒትዎን ለሚከታተለው ሃኪም ይደውሉለት</li> <li>- የአንጀት ምርመራ /ኮሎኖስኮፒ እንዳለበት እና ቀኑን ለህክምና አቅራቢ ይንገሩት</li> <li>- ለስኳር መድሃኒትዎ የተለየ መመሪያ ካለ ይጠይቁ።</li> </ul> </li> </ul> | <p>ቀን</p> <ul style="list-style-type: none"> <li><b>ለቁርስ:-</b> ቀለል ያሉ ክራከሮችን (የተጠበሰ ዳቦ ጥራጥሬ የሌለበት)፣ የበሰለ ሴሪያል(የፋሪና ወይም የስንዴ እልበት፣ግራትሰ)፣ ቀዝቃዛ ሴሪያል(ኮርን ፍሌክስ፣የሩዝ ኮቸሮ)፣ ነጭ ሩዝ፣ኖዳል ወይም ፓስታ</li> <li><b>ጠዋት ላይ</b> :-የሆድዕቃ ማፅጃ ፈሳሽን(ሙቪፕሬፕ)ን ያዘጋጁ።"አንድ እጅ(ፖውች) ኤ" እና "አንድ እጅ (ፓውች) ቢ" በሙሉ ወደ እቃው ይገልብጡ። እቃው ላይ ባለው መስመር ድረስ ለብ ያለ ውሃ ይጨምሩ። ከዛ ማቀዝቀዣ ፍራጅ ስር ያስቀምጡት ፡</li> <li><b>ከ 12 ሰዓት(ቀትር) በኋላ</b>:-ምንም ዓይነት ጠጣር ምግብ አይመገቡ።በቀሪው የቀንዎ ጊዜ ጥርት ያለ ፈሳሽ እንደ ጭማቂ(ቱፋህ፣ወይን ወይም ክራንቤሪ) አጥሚት፣ጥቁር ቡና ወይም ሻይ፣የስፖርት መጠጦች፣ለስለሳ ወይም ጄላቲን(አረንጓዴ ወይም ቢጫ ብቻ )</li> <li><b>7 p.m ላይ</b> :- በየ 15 ደቂቃው 8 አውንስ የሆድዕቃ ማፅጃ/ሙቪ ፕሬፕን ይጠጡ። ከዛ በኋላ 32 አውንስ ጥርት ያለ ፈሳሽ ይጠጡ። ይህ የግድ ተቆማጥ መፍጠር አለበት።ሁለተኛውን እቃ ያዘጋጁ።ፍራጅ ውስጥም ያኑሩት።</li> </ul> | <p>ቀን</p> <ul style="list-style-type: none"> <li>ምንም ዓይነት ጠጣር ምግብ አይብሉ።</li> <li><b>ከቀጠሮ 5 ሰዓታት ቀደም ብሎ</b> :- የሆድዕቃ ማፅጃ ፈሳሽ(ሙቪፕሬፕ)ን 2ኛ እቃ መጠጣት ይጀምሩ። በየ 15 ደቂቃ ልዩነት 8 አውንስ ይጠጡ። ሙቪፕሬፕን ጠጥተው ከጨረሱ በኋላ ቢያንስ 16 አውንስ የሚሆን ጥርት ያለ ፈሳሽ ይጠጡ።</li> <li>የሆድዕቃ ማፅጃ(ሙቪፕሬፕን) ከጠጡ ከ 1 ሰዓት በፊት ወይም በኋላ ጠዋት ላይ የሚወስዷቸውን መድኃኒቶች በውሃ ይዋጧቸው።</li> <li>ለቀጠሮ 2 ሰዓት እስኪቀረው ድረስ ጥርት ያለ ፈሳሽ መጠጣትዎን ይቀጥሉ።</li> <li><b>ቀጠሮ 2 ሰዓት ከቀረው ጊዜ ጀምሮ</b> :-ምንም ዓይነት ነገር በአፍዎ መውሰድ የለብዎትም።</li> </ul> <p><b>ወደ ቀጠሮ ሲመጡ ማምጣት ያለብዎት ነገሮች:-</b></p> <ul style="list-style-type: none"> <li>አዋቂ የቤተሰብ አባል ወይም ጓደኛ</li> <li>በአሁኑ ወቅት የሚወሰዷቸውን ሁሉንም ዓይነት መድሃኒቶች</li> <li>መታወቂያ፣የኢንሹራንስ ካርድ እና አስፈላጊ ከሆነ የወጭ መጋራት ክፍያ (ኮፔይ)</li> <li>ከአንጀት ምርመራው በፊት ሁሉንም ጌጣጌጦች ማውለቅ ይኖርበዎታል።ከተቻለ ውድ ውድ እቃዎችንም ቤት ያስቀምጡ፤አለብዎት የቤተሰብ አባል ወይም ጓደኛዎ እንዲያስቀምጥልዎ ያድርጉ።</li> </ul> |

## የህክምና ቦታ እና የመኪና ማቆሚያ

### ሃርበርቪው ህክምና ማእከል

የኢንዱስትሪ ክፍል  
Maleng Building, 8th Floor  
410 Ninth Ave., Seattle, WA 98104

**206.744.3519**

የመኪና ማቆሚያ 7 ቀን በሳምንት እና 24 ሰዓት በቀን በ 9ኛውና ጀፈርሰን ጋራጅ(P2) ከጄምስ እና ቴሪ ተቃራኒ/ትይዩ

### የዋሽንግተን ዩኒቨርሲቲ ህክምና ማዕከል

የምግብ አፈጫጨት ጤና ማዕከል  
Surgery Pavilion, 3rd Floor  
1959 N.E. Pacific St., Seattle, WA 98195

**206.598.4377**

ከ N.E. Pacific St., ወደ ድንገተኛ ህክምና ክፍል የሚወስደውን አማላካች ይከተሉ። ከላይኛው ጠቋሚ, ወደ Surgery Pavilion Parking Garage በኩል ወደ ግራ ይታጠፉ። መኪናዎችን ካቆሙ በኋላ ወደ 3 ኛው ፎቅ ለመሄድ ሊጭቱን/አሳንሳሩን ይጠቀሙ። ከዛ ወደ ምግብ አፈጫጨት ጤና ማዕከል ለመሄድ ወደ ቀኝ ይታጠፉ።

### ኖርዝ ዌስት ሆስፒታል

1550 N. 115th St., Seattle, WA 98133

**206.368.1770**

የኢንዱስትሪ ምርመራ ክፍል የሚገኘው ከዋናው ሆስፒታል በመጀመሪያው ፎቅ ላይ ስታር ባክስን ተሻግሮ ነው።

### ኢስትላይድ ስፔሻሊስቶች ማዕከል

3100 Northup Way, Bellevue, WA 98004

**425.646.7777**

Free parking on site.

### ሲያትል ካንሰር ኬር አሊያንስ

825 Eastlake Ave. E., Seattle, WA 98109-1023

**206.606.1329**

ምርመራ ክፍሉ የሚገኘው በክሊኒኩ 2ኛ ፎቅ ላይ ነው።

## ልዩ መመሪያዎች

- የስኳር ህመም መድኃኒት የሚወስዱ ከሆነ ፡ የአንጀት ምርመራ /ኮሎኖስኮፒ በሚደረግበት ቀን እና ከዛ ግዜ ቀደም ብሎ ላለው ቀን መድኃኒትዎ መስተካከል ይኖርበታል። በነኝህ 2 ቀናት ውስጥ ምን ማድረግ እንዳለበዎት የስኳር ህመም ህክምና አቅራቢዎች ጋር ይወያዩ።
- የሆድ ድርቀት ካለበዎት አለያም የህመም ማስታገሻ በየቀኑ የሚወስዱ ከሆነ
  - 1 ጠርመራ ማግኘት/የም ሲትሬት ከመድኃኒት መደብር ይግዙ።
  - ለምርመራው 2 ሙሉ ቀናት ሲቀሩት ማታ 8 p.m ላይ ሙሉ ብልቃጡን ይጠጡ።

## ኮሎኖስኮፒ ምንድነው?

ኮሎኖስኮፒ ዶክተርዎ የትልቁ አንጀት ውስጣዊ ንጣፍን እንዲያይ የሚያስችል ምርመራ ነው።

## ምን ሊከሰት እንደሚችል መጠበቅ ያለበዎት

በክርንዎ ደም ስር(ቪይን) ላይ በተተከለው የምግብ መርፌ አማካኝነት ሰመመን ውስጥ የሚከት መድኃኒት ይሰጠዎታል። አብዛኛዎቹ ታካሚዎች በዚህ ወቅት ይተኛሉ። እርስዎ ካልተኙ እና የምችት ስሜት ካልተሰማዎት እባክዎን ይደውሉ። ምችት እንዲሰማዎት የተቻለንን እናደርጋለን።

ዶክተርዎ ተጣጣፊ የሆነ ቱቦ በፊንጢጣዎ ውስጥ ይከታል። የቱቦው መጠን የእጅ ጣትን ያክላል። ዶክተርዎ የህብረ-ህዋስ ናሙና ሊወስድ ይችላል፤ አለያም እብጠቶችን ቆርጦ ሊያስወግድ ይችላል።

የአንጀት ምርመራው /ኮሎኖስኮፒ ከ 30-45 ደቂቃ ይፈጃል። ነገር ግን ለዝግጅት እና ከህክምና በኋላ ለማገገም ሲባል ቀጠሮዎ ረጅም ግዜ ሊወስድ ይችላል።

ከዛ በኋላ የእርስዎ ዶ/ር ስለ ውጤቱ ሊነግርዎት ይችላል። ውጤቱ ወዲያውኑ ካልደረሰ መቸ ሊጠናቀቅ እና ሊደርስዎት እንደሚችል እናሳውቀዎታለን።

### ጥያቄዎች?

ጥያቄዎቻችሁ በጣም ጠቃሚዎች ናቸው ። እባክዎን ጥያቄ ወይም ሀሳብ ካለዎት ይደውሉ። በስራ ቀናት ከ 8 a.m እስከ 5 p.m ፤ የኮሎኖስኮፒ ምርመራ ቀጠሮ ወደታያዘበት ክሊኒክ ይደውሉ። ከስራ ቀናት/ሰዓት ውጭ፤ ቅዳሜና እሁድ ወይም በበዓላት ቀናት ሃርበርቪው የሚገኘውን የማህበረሰብ ክብካቤን በ 206.744.2500 ይደውሉ።

# Colonoscopy

## How to prepare with MoviPrep

Your colonoscopy is scheduled for \_\_\_\_\_, \_\_\_\_\_, at \_\_\_\_\_  
DAY DATE TIME

| 7 Days Before  | 5 Days Before  | 1 Day Before   | On Procedure Day  |
|--|--|--|---|
| Date:  | Date:  | Date:  | Date:   |
| <ul style="list-style-type: none"> <li>• <b>If you take blood-thinning medicines</b> such as Coumadin (warfarin) or Plavix (clopidogrel):                             <ul style="list-style-type: none"> <li>- Call the doctor who prescribed these medicines.</li> <li>- Tell the doctor you are having a colonoscopy, and the date of your procedure.</li> <li>- Ask for special instructions for your blood-thinning medicines.</li> </ul> </li> <li>• <b>Cancellations:</b> If you need to reschedule or cancel your test, call 206._____.</li> <li>• <b>Discharge and transportation:</b> <ul style="list-style-type: none"> <li>- An adult family member or friend <b>must</b> pick you up when you are ready to go home. You cannot drive, take a bus or taxi, or use a service such as Hopelink or Paratransit by yourself after your test.</li> <li>- <b>Your test will be cancelled if you do not have an adult family member or friend to take you home.</b></li> <li>- We strongly advise that a responsible adult stay with you for the rest of the day.</li> </ul> </li> </ul> | <ul style="list-style-type: none"> <li>• <b>Stop</b> eating foods that contain nuts or seeds.</li> <li>• <b>Stop</b> taking iron and any multivitamin that contains iron.</li> <li>• <b>Fill</b> your MoviPrep prescription.</li> <li>• <b>Read</b> this handout carefully.</li> <li>• <b>If you have diabetes:</b> <ul style="list-style-type: none"> <li>- Call the provider who manages your diabetes medicines.</li> <li>- Tell the provider that you are having a colonoscopy and the date of your procedure.</li> <li>- Ask for special instructions for your diabetes medicines.</li> </ul> </li> </ul> | <ul style="list-style-type: none"> <li>• <b>For breakfast:</b> You can eat plain crackers (saltines or Melba toast, no seeds), cooked cereals (farina, cream of wheat, grits), cold cereal (corn flakes, puffed rice), white rice, noodles, or pasta.</li> <li>• <b>In the morning:</b> Prepare the MoviPrep solution. Empty one "Pouch A" and one "Pouch B" into the container. Add lukewarm water to the top line of the container. Store in the refrigerator.</li> <li>• <b>After 12 noon:</b> Do not eat any solid food. For the rest of the day, drink clear liquids such as juice (apple, grape, or cranberry), broth, black coffee or tea, sports drinks, soda, or gelatin (green or yellow only).</li> <li>• <b>At 7 p.m.:</b> Drink 8 ounces of MoviPrep every 15 minutes. Then drink at least 32 ounces of clear liquid. It should cause diarrhea. Prepare the 2nd container. Store it in the refrigerator.</li> </ul> | <ul style="list-style-type: none"> <li>• <b>Do not eat any solid food.</b></li> <li>• <b>5 hours before your appointment:</b> Start drinking the 2nd container of MoviPrep. Drink 8 ounces every 15 minutes. After you have finished the MoviPrep, drink at least 16 ounces of a clear liquid.</li> <li>• 1 hour before or after drinking the MoviPrep, take your morning medicines with water.</li> <li>• Keep drinking clear liquids until 2 hours before your appointment.</li> <li>• <b>Starting 2 hours before your appointment:</b> Do not take anything by mouth.</li> </ul> <p><b>What to bring to your appointment:</b></p> <ul style="list-style-type: none"> <li>• An adult family member or friend</li> <li>• A list of all medicines you currently take</li> <li>• Photo ID, insurance card, and co-pay, if needed</li> <li>• You will need to remove all jewelry and piercings before your colonoscopy. If you can, leave all valuables at home, or give them to your family member or friend for safekeeping.</li> </ul> |

## Locations and Parking

### Harborview Medical Center

GI Endoscopy Suite  
Maleng Building, 8th Floor  
410 Ninth Ave., Seattle, WA 98104  
**206.744.3519**

Parking 24 hours a day, 7 days a week in the Ninth and Jefferson Garage (P2) off James and Terry.

### University of Washington Medical Center

Digestive Health Center  
Surgery Pavilion, 3rd Floor  
1959 N.E. Pacific St., Seattle, WA 98195  
**206.598.4377**

From N.E. Pacific St., follow signs to the Emergency Room. At the stop sign, turn left into the Surgery Pavilion Parking Garage. After parking, take the elevator to the 3rd floor. Turn right into the Digestive Health Center.

### Northwest Hospital

1550 N. 115th St., Seattle, WA 98133  
**206.368.1770**

The Endoscopy Procedure Suite is on the 1st floor of the main hospital, across from Starbucks.

### Eastside Specialty Center

3100 Northup Way, Bellevue, WA 98004  
**425.646.7777**

Free parking on site.

### Seattle Cancer Care Alliance

825 Eastlake Ave. E., Seattle, WA 98109-1023  
**206.606.1329**

The Procedure Suite is on the 2nd floor of the clinic.

## Special Instructions

- **If you take medicines for diabetes:** Your medicines will need to be adjusted for the day before and the day of your colonoscopy. Talk with your diabetes care provider about what to do on these 2 days.
- **If you have severe constipation or are taking pain medicine every day:**
  - Buy a bottle of Magnesium Citrate at a drugstore.
  - 2 days before your procedure, at 8 p.m., drink the full bottle.

## What is a colonoscopy?

A *colonoscopy* allows your doctor to look at the lining of your large intestine.

## What to Expect

You will receive a *sedative* (medicine to help you relax) through an *intravenous line* (IV) placed in your vein. Most patients fall asleep. If you do not fall asleep and feel some discomfort, please tell us. We will do our best to help you be comfortable.

Your doctor will place a flexible tube into your rectum. The tube is about as wide as a finger. Your doctor may also take *biopsies* (tissue samples) or remove *polyps* (growths).

The colonoscopy takes about 30 to 45 minutes. Your appointment may last longer to allow time to prepare and recover.

Afterward, your doctor may talk with you about the results. If your results are not ready right away, we will tell you when you can expect to receive them.

## Questions?

Your questions are important. Please call if you have questions or concerns. Weekdays between 8 a.m. and 5 p.m., call the clinic where your colonoscopy is scheduled. After hours and on weekends and holidays, call the Community Care Line at Harborview, 206.744.2500.